

ZOETIS INC.

333 PORTAGE STREET, KALAMAZOO, MI, 49007

Telephone: 269-359-4414

Customer Service: 888-963-8471

Website: www.zoetis.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

FluSure XP®/FarrowSure® GOLD B

Zoetis

Parvovirus-Swine Influenza Vaccine, H1N1 & H1N2 & H3N2

Killed Virus

Erysipelothrix Rhusiopathiae-Leptospira Bratislava-Canicola-Grippotyphosa-Hardjo-Icterohaemorrhagiae-Pomona Bacterin

For use in swine only

For veterinary use only

For vaccination of healthy breeding swine, including pregnant sows and gilts, as an aid in preventing respiratory disease caused by swine influenza A viruses H1N1, H1N2 and H3N2, reproductive failure caused by PPV, and disease caused by *Erysipelothrix rhusiopathiae*, *Leptospira bratislava*, *L. canicola*, *L. grippotyphosa*, *L. hardjo*, *L. icterohaemorrhagiae* and *L. pomona*. An 18-week duration of immunity following vaccination has been demonstrated against erysipelas. This product contains the following Influenza A Virus isolates: A/Swine/Iowa/110600/2000 (H1N1); A/Swine/Oklahoma/0726H/2008 (H1N2); A/Swine/North Carolina/394/2012 (H3N2); and A/Swine/Minnesota/872/2012 (H3N2).

Directions: Aseptically rehydrate the freeze-dried vaccine with the liquid component provided, shake well, and administer 2 mL intramuscularly. Healthy swine should receive 2 doses 3-5 weeks apart with the second dose administered 2-4 weeks prior to breeding. Healthy gilts, however, should receive the second dose as near as possible to 14 days prior to breeding. Revaccination with a single dose is recommended prior to subsequent breedings. Boars should be revaccinated semiannually.

Precautions: Store at 2°-7°C. Do not freeze. Use entire contents when first opened or rehydrated. Sterilized syringes and needles should be used to administer this vaccine. Do not vaccinate within 21 days before slaughter. Contains gentamicin as preservative. Temporary swelling at the injection site may be observed. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471 (USA), (800) 461-0917 (Canada).

U.S. Veterinary License No. 190

Zoetis Inc., Kalamazoo, MI 49007, USA

	Code	
50-dose vial of vaccine, rehydrate to 100 mL	49F6.2B	50050700
100-mL vial of liquid component for use as diluent		

CPN: 3690249.5