

ARKO LABORATORIES, LTD.

P.O. BOX 400, 1336 MAIN STREET, JEWELL, IA, 50130

Telephone: 515-827-5648

Toll-Free: 800-714-2756

Fax: 515-827-5648

Website: www.arkolabs.com

Email: arko@arkolabs.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

EDEMA VAC 100

Arko

ESCHERICHIA COLI VACCINE

Avirulent Live Culture

INDICATIONS: The product is recommended as an aid in the prevention of edema disease caused by F18⁺ E. Coli in swine.

DIRECTIONS AND DOSAGE: Add 20 mL of water, shake well. Combine with 80 mLs of water and promptly administer 1 mL orally per pig. Pigs should be at least 18 days of age or older when vaccinated.

PRECAUTIONS: Store at not over 45°F (7°C). Use entire contents when first opened. Burn this container and all unused contents. Do not vaccinate within 21 days before slaughter. Anaphylactic reactions occur. Antidote: administer epinephrine.

100 Doses

U.S. VETERINARY LICENSE NO. 337

ARKO LABORATORIES LTD., JEWELL, IOWA

CPN: 1123008.1

ARKO LABORATORIES, LTD.

P.O. BOX 400, 1336 MAIN STREET, JEWELL, IA, 50130

Telephone: 515-827-5648

Toll-Free: 800-714-2756

Fax: 515-827-5648

Website: www.arkolabs.com

Email: arko@arkolabs.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

EDEMA VAC 500

Arko

ESCHERICHIA COLI VACCINE

Avirulent Live Culture

INDICATIONS: The product is recommended as an aid in the prevention of edema disease caused by F18⁺ E. Coli in swine.

DIRECTIONS AND DOSAGE: Add 20 mLs. of water, shake well. Combine with 480 mLs. of water and promptly administer 1 mL. orally per pig. Pigs should be at least 18 days of age or older when vaccinated.

PRECAUTIONS: Store at not over 45°F (7°C). Use entire contents when first opened. Burn this container and all unused contents. Do not vaccinate within 21 days before slaughter. Anaphylactic reactions occur. Antidote: administer epinephrine.

500 Doses

U.S. VETERINARY LICENSE NO. 337

ARKO LABORATORIES LTD., JEWELL, IOWA

CPN: 1123015.0