

Porcilis™ Ileitis

LAWSONIA INTRACELLULARIS VACCINE BACTERIN

For use in healthy swine as an aid in the control of ileitis caused by *Lawsonia intracellularis*, an aid in the reduction of colonization by *Lawsonia* and an aid in the reduction of duration of fecal shedding. Duration of immunity for at least 20 weeks has been demonstrated.

Advantages

Porcilis™ Ileitis offers long-lasting protection for ileitis control.

- Industry-leading 20-week (5 months) duration of immunity (DOI)
 - With a 20-week DOI, vaccination with Porcilis Ileitis provides 13 more weeks of protection than the water-administered vaccine on the market
- One-dose, ready-to-use injectable vaccine
 - As the only injectable vaccine available for ileitis, Porcilis Ileitis allows for vaccination of pigs at 3 weeks of age or older – an age when they're easier to handle
 - A single-bottle vaccine means no mixing, combining or risk of contamination from the process
 - No worrying about removing feed-grade antibiotics during vaccination



New peel-off sticker enables producers to easily record when vaccinations are given.



Code: 126464 100 mL
For piglets: 50 doses – 2 mL dose

Code: 144648 500 mL
For piglets: 250 doses – 2 mL dose

Porcilis™ Ileitis

LAWSONIA INTRACELLULARIS VACCINE BACTERIN

For use in healthy swine as an aid in the control of ileitis caused by *Lawsonia intracellularis*, an aid in the reduction of colonization by *Lawsonia* and an aid in the reduction of duration of fecal shedding. Duration of immunity for at least 20 weeks has been demonstrated.

ADMINISTRATION AND DOSAGE:

Shake well. Warm the vaccine to room temperature immediately prior to administration. Using aseptic technique, inject 2 mL once intramuscularly (IM) into pigs at 3 weeks of age or older.

CAUTIONS:

Store in the dark at 2° to 7°C (35° to 45°F). Do not freeze. Use entire contents when first opened; do not save partial contents. Burn the container and all unused product. Use only in healthy swine. Do not vaccinate within 21 days before slaughter. A significant number of pigs may experience transient swelling at the site of injection. The swelling should resolve without treatment in less than 20 days. **Immediate systemic reactions have been observed in animals administered this product. Minimize pig handling and ensure proper ventilation prior to starting vaccination.** If an allergic reaction occurs, treat with epinephrine. Caution should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. Contact a physician immediately.

PRESERVATIVE:

Gentamicin and thimerosal

FOR ANIMAL USE ONLY

READ AND FOLLOW LABEL DIRECTIONS

U.S. Veterinary License No. 165A

FOR VETERINARY USE ONLY