

General Lube no pump



A viscous bland lubricant which is non-spermicidal for gynecological and rectal exams.

Size	Reorder Number	Lbs/Case	Case Pack
Gallon	LU020PCNP	35 lbs	4

NDC# 58829-150-01



DIRECTIONS FOR USE: A non-sterile non-spermicidal all purpose lubricating jelly for use when performing gynecological procedures and rectal exams.

The bland lubricating qualities aid in minimizing irritation to both the operator's skin and membranes of the animal.

Apply liberally as needed.

General Lube

All Purpose Non-Spermicidal
Lubricating Jelly

INGREDIENTS: Purified water, propylene glycol, sodium carboxymethylcellulose, methyl and propyl parahydroxybenzoate.

STORAGE: Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.



Priority Care is a registered trademark of First Priority, Inc.

For Animal Use Only • Keep Out of Reach of Children

Net Contents: One Gallon (3.785 L)

LOT NO.

Reorder No. LU020PC

Rev. 02-07

Manufactured By:
FIRST PRIORITY, INC., Elgin, IL 60123-1146

DURVET, INC.

100 S.E. MAGELLAN DRIVE, BLUE SPRINGS, MO, 64014

Telephone: 816-229-9101

Toll-Free: 800-821-5570

Fax: 816-224-3080

Website: www.durvet.com

Email: info@durvet.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

MAXI-SORB BOLUS

Durvet

Antidiarrheal-Demulcent

Aids in relief of simple diarrhea in horses and cattle.

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS: Each bolus contains: Activated Attapulgite 20%.

INACTIVE INGREDIENTS: Carob Flour, Pectin and Magnesium Trisilicate, in a palatable base.

Store at room temperature between 15° and 30°C (59°-86°F).

Indications: For use as an aid in relief of simple non-infectious diarrhea in horses and cattle.

DOSAGE AND ADMINISTRATION: Administer orally. Give two (2) boluses to adult cattle and horses. Give one (1) bolus to calves and foals.

Repeat treatment at 4 to 6 hour intervals as needed. Discontinue use of product after 3 days. If symptoms persist after using this preparation for 3 days, consult a veterinarian.

Manufactured for: **DURVET, INC.**, Blue Springs, Missouri 64014

www.durvet.com

NET CONTENTS: NET WEIGHT PER BOLUS: NDC

50 BOLUSES 17.9g - 18.9g

30798-016-69

T-7667-02

Rev. 05-11

CPN: 1084114.4

