

**INDICATIONS:** GEN-GARD Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin, and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

**DOSE:** Colibacillosis - 1 level scoop per 240 gallons of drinking water for 3 consecutive days, 0.5 mg/lb/day. For proportioner use, add 1 level scoop to 2 gallons stock solution and dispense at the rate of 1 oz./gal. drinking water (25 mg/gal.).

Swine Dysentery - 1 level scoop per 120 gallons of drinking water for 3 consecutive days, 1.0 mg/lb/day. For proportioner use, add 1 level scoop to 1 gallon stock solution and dispense at the rate of 1 oz./gal. drinking water (50 mg/gal.).

**FOR USE IN ANIMALS ONLY**  
**KEEP OUT OF REACH OF CHILDREN**



G-7473-03

Rev. 07-16

© REGISTERED TRADEMARK OF AGRI LABORATORIES, LTD.

MFD. FOR AGRI LABORATORIES, LTD.  
St. Joseph, MO 64503



NDC 57561-474-03

**GEN-GARD®**

Gentamicin Sulfate  
Soluble Powder

**ANTIBACTERIAL**

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

Contains 120 grams of  
Gentamicin per jar

ANADA 200-185, Approved by FDA

**NET WEIGHT:**  
**360 g (12.7 oz.)**



 **AgriLabs®**  
Making A Healthy Difference®

**EACH G CONTAINS:** Gentamicin sulfate equivalent to 333.33 mg gentamicin base.

**RESIDUE WARNING:** Swine treated with the recommended dose of **GEN-GARD Soluble Powder** must not be slaughtered for food for at least ten (10) days following the last treatment.

Do not store or offer medicated water in rusty metal containers.

**Read accompanying directions carefully.**

**Store between 2° and 30°C (36° and 86°F).**

**For use in swine drinking water only.**

**Restricted drug (California)-**  
**Use only as directed.**

Lot No.

Exp. Date