

ORAL-PRO™

Sodium Salicylate Concentrate 48.6% w/v



FEATURES AND BENEFITS

- Aids in reducing fever caused by viral and bacterial infections.^{1,2}
- Aids in reducing fever caused by heat stress from transporting.
- Offers water solubility in hard water, and is also stable in milk.³
- Offers stress relief from vaccination and processing.
- Manufactured under cGMP standards, guaranteeing consistent quality.
- Calves: A supportive treatment of pyrexia in acute respiratory disease and in combination with appropriate (anti-infective) therapy, if necessary.

¹ Lees, P. and S.A. May, 1992. Inflammation and anti-inflammatory drugs. Pages 843–863 in: Bovine Medicine. A.H. Andrews, R.W. Blowey, H. Boyd, and R.G. Eddy, ed. Oxford Blackwell Scientific Publications, Cambridge, MA.

² Osol, A., R. Pratt, and A.R. Gennaro, 1973. The United States Dispensatory, 27th ed. J.B. Lippincott Co., Philadelphia, PA.

³ Budavari, S., M.J. O'Neil, A. Smith, and P.E. Heckelman, 1989. The Merck Index, 11th ed. Merck & Co., Rahway, NJ.

INDICATIONS

For use in the drinking water of calves, poultry and swine as an aid in reducing pain, fever and inflammation.

FOR USE IN LIVESTOCK ONLY

REORDER NO: 21004

NDC 51072-038-01

Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256 • www.aurorapharmaceutical.com

12/2011 IN 50-1108 REV 01
ss000003


MANUFACTURED IN THE USA


aurora
PHARMACEUTICAL[®]



Sodium Salicylate Concentrate 48.6% w/v Solution

APPLICATION

Administer 4 ml per 100 lbs. body weight (40 mg/kg body weight Sodium Salicylate) to calves once daily for 1–3 days. Administer orally in drinking water or milk (replacer). Zero day withdrawal period for meat and offal from calves. Do not use in cows producing milk for human consumption.

INDICATIONS: For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

CALF LABEL CLAIMS

Indications and Use, Calves: Suppressive treatment of gross respiratory disease associated with appropriate (e.g., anti-infective) therapy if necessary.

Contraindications, Calves: Do not use Sodium Salicylate in neonates or calves less than 1 week of age.

Dosage, Calves: 40 mg Sodium Salicylate per kg body weight once daily, for 1–3 days. Administer orally in drinking water or milk (replacer).

Days Recommended withdrawal period for all food animals following use at "usual dosages": 24 hours.

ACTIVE INGREDIENT: Sodium Salicylate.....48.6% w/v

DIRECTIONS FOR USE:

For Analgesic and Antipyretic Use:

Water: Add 8 ounces (236 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

Livestock Tank Use: Add 8 ounces (236 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

Following Days: Water Product Use: Add 10 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Medicoder Tank Use: Add 10 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Prepare fresh solutions daily. Repeat as necessary.

WARNING: Do not administer concentrated solution by direct oral administration — gastritis and/or diarrhea may occur. Do not use in animals less than 1 week of age.

Store at controlled room temperature between 15°–30° C (59°–86° F).

CAUTION: Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place container in warm water. Gentle shaking will restore consistency or product.

Gradual darkening will not affect product stability.

REORDER NO: 21004



NDC 51072-038-01

INDICACIONES: Para uso en el agua potable de las aves de corral y cerdos como ayuda para reducir el dolor, fiebre e inflamación.

TENEREAS DE LA ETIQUETA

Indicaciones de la Terapéutica: Tratamiento de la lesión de enfermedad respiratoria aguda, incluyendo la bronquitis (por ejemplo, anti-infeccioso), la terapia, si es necesario.

Contraindicaciones, Terapéuticas: No utilizar salicilato de sodio en los recién nacidos o terneras de menos de 2 semanas de edad.

Dosis, Terapéuticas: 40 mg de salicilato sodico por kg de peso corporal una vez al día, durante 1–3 días. Administrar en agua potable o leche (sustituto). De suspensión recomendado periodo para todos los animales de alimento después de su uso en "dosis habituales"; la carne de 24 horas.

INGREDIENTES ACTIVOS: Sodio salicilato.....48.6% p/b

MODO DE EMPLEO: Para el Uso de Analgésicos y Antipiréticos

Uso del Difusor de Agua: Agregar 8 onzas (236 ml) de concentrado de salicilato de sodio 48.6% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetiva de 11.3 mg/lb (25 mg/kg) de peso corporal al dia.

Uso de Tanque Ganadero: Agregar 8 onzas (236 ml) de concentrado de salicilato de sodio 48.6% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetiva de 11.3 mg/lb (25 mg/kg) de peso corporal al dia.

Para el Uso de Analgésicos y Antipiréticos:

Uso del Difusor de Agua: Agregar 10 onzas (473 ml) de concentrado de salicilato de sodio 48.6% para hacer un galón de solución madra y administrar a través de un medicador medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetiva de 13.6 mg/lb (30 mg/kg) de peso corporal al dia.

Uso del Tanque Ganadero: Agregar 10 onzas (473 ml) de concentrado de salicilato de sodio 48.6% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetiva de 13.6 mg/lb (30 mg/kg) de peso corporal al dia.

Los Días Siguientes: Uso del Difusor de Agua: Agregar 10 onzas (473 ml) de concentrado de salicilato de sodio 48.6% para hacer un galón de solución madra y administrar a través de un medicador medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetiva de 13.6 mg/lb (30 mg/kg) de peso corporal al dia.

Uso del Tanque Ganadero: Agregar 10 onzas (473 ml) de concentrado de salicilato de sodio 48.6% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetiva de 13.6 mg/lb (30 mg/kg) de peso corporal al dia.

Preparar soluciones frescas diariamente. Repita según sea necesario.

ADVERTENCIA: No administrar la solución concentrada por vía oral en directo — irritación gastrintestinal o una sobredosis pueden ocurrir. No utilizar en lechones de menos de 4 semanas de edad.

Consejo: Mantenga el recipiente cerrado cuando no esté en uso.

El producto puede solidificarse a temperaturas de almacenamiento en frío. Coloque el recipiente en agua tibia para derretirlo. No caliente el recipiente directamente en la solución, o en lugar con agua tibia. Inverta suavemente el recipiente para asegurar la uniformidad del producto.

Gradualmente el efecto no afectará a la estabilidad del producto.

Oscurecimiento gradual no afectará a la estabilidad del producto.

ORAL-PRO™ Sodium Salicylate Concentrate 48.6% w/v

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:
1 Gallon (3.79 Liters)



ABOUT AURORA PHARMACEUTICAL

Aurora is devoted to improved formulations and delivery systems that are consumer friendly and which utilize state-of-the-art ingredient "green" technologies. We are committed to manufacturing products with top-quality ingredients that increase the stability, shelf life and efficacy of our products while minimizing harmful ingredients and environmental contamination.

Aurora Pharmaceutical, LLC

NORTHFIELD, MINNESOTA 55057

888-215-1256 • www.aurorapharmaceutical.com



MANUFACTURED IN THE USA