

# HUVEPHARMA, INC.

525 WESTPARK DR., SUITE 230, PEACHTREE CITY, GA, 30269

Telephone: 770-486-7212

Toll-Free: 877-994-4883

Fax: 770-486-7217

Website: [www.huvepharma.com](http://www.huvepharma.com)



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

## TYLOVET® SOLUBLE



### **Huvepharma**

#### **Tylosin Tartrate**

Equivalent to **100 g (3.53 oz)** tylosin base

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**For oral use in chickens, turkeys, swine, and honey bees.**

Macrolide Antibiotic, ANADA 200-473, approved by FDA

#### **GRANULES FOR ORAL SOLUTION**

#### **INDICATIONS**

**Chickens:** As an aid in the treatment of chronic respiratory disease (CRD) associated with *Mycoplasma gallisepticum* sensitive to tylosin in broiler and replacement chickens. For the control of CRD associated with *Mycoplasma gallisepticum* sensitive to tylosin at the time of vaccination or other stress in chickens. For the control of CRD associated with *Mycoplasma synoviae* sensitive to tylosin in broiler chickens.

**Turkeys:** For the reduction in severity of effects of infectious sinusitis associated with *Mycoplasma gallisepticum*.

**Swine:** For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae*. For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* when followed immediately by tylosin phosphate Type A medicated article in feed. For the control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* when followed immediately by tylosin phosphate Type A medicated article in feed.

**Honey Bees:** For the control of American Foulbrood (*Paenibacillus larvae*).

#### **INGREDIENTS**

Tylosin (as tylosin tartrate)	100 g
-------------------------------	-------

#### **DOSAGE AND ADMINISTRATION**

**Chickens and Turkeys:** To assure thorough dissolution, place the **Tylovet** (contents of the pouch) in a mixing container and add one gallon of water (3790 mL) to the material. Mix this concentrated solution with water to make 50 gallons (189 liters) of treated drinking water resulting in 2,000 mg/gallon.

**Swine:** To assure thorough dissolution, place the **Tylovet** (contents of the pouch) in a mixing container and add one gallon of water (3790 mL) to the material. Mix this concentrated solution with water to make 400 gallons (1514 liters) of treated drinking water resulting in 250 mg/gallon.

#### **Mixing Directions for Water Medicating Pump (1:128 inclusion)\***

Species	Desired Concentration in Drinking Water (mg/gallon)	Pouches of Tylovet Soluble	Gallons of Water to Make Stock Solution
Chickens and Turkeys	2,000	10	3.9
Swine	250	1	3.1

\*This table applies only if the water medicating pump is set to deliver 1 ounce of stock solution per gallon of drinking water.

**Honey Bees:** Mix 200 mg tylosin in 20 g confectioners/powdered sugar. Use immediately.

**When directed to mix the product with water, always** add the water to the powder. **Do not pour** the powder into the water. Prepare a fresh **Tylovet** solution every three days.

When mixing and handling tylosin, use protective clothing and impervious gloves.

#### **DIRECTIONS FOR USE:**

**Chickens** should be treated for three days; however, treatment may be administered for one to five days depending upon severity of infection. Treated chickens must consume enough medicated water to provide 50 mg per pound of body weight per day. Only medicated water should be available to the birds.

**Turkeys** should be treated for three days; however, treatment may be administered for two to five days depending upon severity of infection. Treated turkeys must consume enough medicated water to provide 60 mg per pound of body weight per day. Only medicated water should be available to the birds.

**Swine** For the treatment and control of swine dysentery medicate with 250 mg tylosin per gallon in drinking water for 3 to 10 days, depending upon severity of infection. Alternatively, medicate with 250 mg tylosin per gallon in drinking water for 3 to 10 days, followed by 40 to 100 g tylosin per ton of complete feed (Type C medicated feed manufactured from tylosin phosphate Type A medicated article) for 2 to 6 weeks. For control of porcine proliferative enteropathies (PPE, ileitis) medicate with 250 mg tylosin per gallon in drinking water for 3 to 10 days, followed by 40 to 100 g tylosin per ton of complete feed (Type C medicated feed manufactured from tylosin phosphate Type A medicated article) for 2 to 6 weeks. Swine must consume enough medicated water to provide a therapeutic dose. Only medicated water (250 mg tylosin per gallon) should be available while medicating with TYLOVET Soluble.

**Honey Bee colonies** should receive three treatments administered as a dust in confectioners/powdered sugar. The 200 mg dose is applied (dusted) over the top bars of the brood chamber once weekly for 3 weeks.

#### **WARNINGS**

**User Safety Warnings:** Not for Human Use. Keep Out of Reach of Children. Avoid contact with human skin. Exposure to tylosin may cause a rash.

**Residue Warnings:** Chickens must not be slaughtered for food within 24 hours after treatment. Turkeys must not be slaughtered for food within five days after treatment. Swine must not be slaughtered for food within 48 hours after treatment. Do not use in layers producing eggs for human consumption. Honey bees: The drug should be fed early in the spring or fall and consumed by the bees before the main honey flow begins, to avoid contamination of production honey. Complete treatments at least 4 weeks prior to main honey flow.

#### **How supplied:**

Tylovet® Soluble is supplied in pouches of laminated aluminum foil, 100 g of tylosin activity per pouch.

**Storage:** Store at or below 25 °C (77 °F). Excursions Permitted to 40 °C (104 °F). Avoid moisture.

**Restricted Drug (California) - Use Only as Directed.**

**ANADA # 200-473, Approved by FDA**

**To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765.**

**Manufactured for:** Huvepharma Inc. Peachtree City, GA 30269, Made in Bulgaria

Tylovet® is a registered trademark of Huvepharma AD

Product 2220

Code: LB444v3-Tyl100-USA1405

**CPN:** 1309009.2