

Serial No.:
Exp. Date:

Indications: For vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention of porcine proliferative enteropathy (Ileitis) caused by *Lawsonia intracellularis*.

Directions: Rehydrate the vaccine by adding the accompanying diluent to the vaccine vial. Mix well and use immediately.

Dosage: Administer one 2 mL dose per pig, either orally or via the drinking water. See insert for complete directions.

Precautions: Store refrigerated at 35–45°F (2–7°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before

**Lawsonia
Intracellularis Vaccine**
Avirulent Live Culture
Enterisol® Ileitis

For use in swine only

50 Doses/100 mL

slaughter. Burn container and all unused contents. Do not use in pregnant swine or breeding boars. Do not use concurrently with antibiotics. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation and reduced product efficacy. Anaphylactic reactions may occur. Antidote: Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124
Code 130-421



**Boehringer
Ingelheim**

30402-01

Serial No.:
Exp. Date:

Indications: For vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the prevention of porcine proliferative enteropathy (Ileitis) caused by *Lawsonia intracellularis*.

Directions: Rehydrate the vaccine by adding the accompanying diluent to the vaccine vial. Mix well and use immediately.

Dosage: Administer one 1 mL dose per pig, either orally or via the drinking water. See insert for complete directions.

Precautions: Store refrigerated at 35–45°F (2–7°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before

**Lawsonia
Intracellularis Vaccine**
Avirulent Live Culture
Enterisol® Ileitis

For use in swine only

100 Doses/100 mL

slaughter. Burn container and all unused contents. Do not use in pregnant swine or breeding boars. Do not use concurrently with antibiotics. All materials used in administering this vaccine must be free of antimicrobial or disinfectant residue to prevent inactivation and reduced product efficacy. Anaphylactic reactions may occur. Antidote: Epinephrine.

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506
U.S. Veterinary License No. 124
Code 130-441



**Boehringer
Ingelheim**

30403-02