

Ingelvac MycoFLEX®

Mycoplasma hyopneumoniae bacterin
Porcine Circovirus Vaccine Type 2, Killed Baculovirus Vector

Description

Ingelvac MycoFLEX® is a single-dose mycoplasma vaccine. INGELVAC MYCOFLEX has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against *Mycoplasma hyopneumoniae* (Mhp).

Features

- Effective for the vaccination of healthy swine 3 weeks of age or older against *Mycoplasma hyopneumoniae*
- Single dose, 1 mL
- At least 26 weeks' duration of immunity
- Contains ImpranFLEX®, an effective, aqueous-based (non-oil) polymer adjuvant

Benefits

- Low viscosity, highly syringeable
- Rapid onset of immunity as well as long duration of immunity
- PQA® Plus friendly
- Member of the FLEX Family (nine swine vaccines, multiple combinations); allows you a flexible vaccination scheduling
- NOW with a new convenient 500-dose presentation

Dosage & Administration

Shake well before use. Using aseptic technique, administer a single 1-mL dose by intramuscular injection. **Does not need to be tempered.**

Precautions

Store at 35-45° F (2-7° C).
Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.

Uses

Recommended for the vaccination of healthy, susceptible swine 3 weeks of age or older as an aid in the reduction of enzootic pneumonia of swine caused by *Mycoplasma hyopneumoniae*.

Warning

In case of anaphylactic reaction, epinephrine is symptomatic treatment.

Product Information

Name	Code	Size
INGELVAC	138896	50 dose
MYCOFLEX	138898	250 dose
	138897	500 dose

Highlights

- Convenient single 1-mL dose
- Rapid onset of immunity as well as long duration of immunity
- Flexible vaccination scheduling



Serial No.:
Exp. Date:

This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against *Mycoplasma hyopneumoniae*. Duration of immunity is at least 26 weeks. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Composition: Prepared from chemically inactivated cultures of *M. hyopneumoniae*, combined with an adjuvant to stimulate the immune response.

Directions and dosage: Shake well before use. Using aseptic technique, inject a single 1 mL dose intramuscularly. For advice on revaccination frequency, contact your veterinarian or the manufacturer.



Precautions: Store at 35-46°F (2-8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician. **Caution: Do not use as diluent for live vaccines.**

Boehringer Ingelheim
Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/2775.02

 **Boehringer
Ingelheim**

127303-02

Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd., Duluth, Georgia 30096-4640
Information 888-637-4251
www.bi-animalhealth.com/swine